

## 熊本大学学術リポジトリ

### Kumamoto University Repository System

Title	尿中8-hydroxy-2'-deoxyguanosine測定は急性心筋梗塞再灌流療法後の心事故予測因子となる
Author(s)	永吉, 靖央
Citation	
Issue date	2006-01-18
Type	Thesis or Dissertation
URL	<a href="http://hdl.handle.net/2298/3106">http://hdl.handle.net/2298/3106</a>
Right	

## 12. 文献

1. Schilling WP, Elliott SJ. Ca<sup>2+</sup> signaling mechanisms of vascular endothelial cells and their role in oxidant-induced endothelial cell dysfunction. *Am J Physiol.* 1992;262:H1617-1630.
2. Goldstein JL, Kita T, Brown MS. Defective lipoprotein receptors and atherosclerosis. Lessons from an animal counterpart of familial hypercholesterolemia. *N Engl J Med.* 1983;309:288-296.
3. Steinberg D Oxidative Modification of LDL and Atherogenesis. *Circulation.* 1997;95:1062-1071
4. Zhang R, Shen Z, Nauseef WM, Hazen SL. Defects in leukocyte-mediated initiation of lipid peroxidation in plasma as studied in myeloperoxidase-deficient subjects: systematic identification of multiple endogenous diffusible substrates for myeloperoxidase in plasma. *Blood.* 2002;99:1802-1810.
5. Parthasarathy S, Wieland E, Steinberg D. A role for endothelial cell lipoxygenase in the oxidative modification of low density lipoprotein. *Proc Natl Acad Sci U S A.* 1989;86:1046-1050.
6. Cyrus T, Witztum JL, Rader DJ, Tangirala R, Fazio S, Linton MF, Funk CD. Disruption of the 12/15-lipoxygenase gene diminishes atherosclerosis in apo E-deficient mice. *J Clin Invest.* 1999;103:1597-1604.
7. Taniyama Y, Griending KK. Reactive oxygen species in the vasculature: molecular and cellular mechanisms. *Hypertension.* 2003;42:1075-81.
8. Ku DN, Giddens DP, Zarins CK, Glagov S. Pulsatile flow and atherosclerosis in the human carotid bifurcation. Positive correlation between plaque location and low oscillating shear stress. *Arteriosclerosis.* 1985;5:293-302.

9. Kobayashi S, Inoue N, Ohashi Y, Terashima M, Matsui K, Mori T, Fujita H, Awano K, Kobayashi K, Azumi H, Ejiri J, Hirata K, Kawashima S, Hayashi Y, Yokozaki H, Itoh H, Yokoyama M. Interaction of oxidative stress and inflammatory response in coronary plaque instability: important role of C-reactive protein. *Arterioscler Thromb Vasc Biol.* 2003;23:1398–1404.
10. Fuster V, Stein B, Ambrose JA, Badimon L, Badimon JJ, Chesebro JH. Atherosclerotic plaque rupture and thrombosis. Evolving concepts. *Circulation.* 1990;82:II47–II59.
11. Libby P. Molecular bases of the acute coronary syndromes. *Circulation.* 1995;91:2844–2850.
12. van der Wal AC, Becker AE, van der Loos CM, Das PK. Site of intimal rupture or erosion of thrombosed coronary atherosclerotic plaques is characterized by an inflammatory process irrespective of the dominant plaque morphology. *Circulation.* 1994;89:36–44.
13. Rajagopalan S, Meng XP, Ramasamy S, Harrison DG, Galis ZS. Reactive oxygen species produced by macrophage-derived foam cells regulate the activity of vascular matrix metalloproteinases in vitro. Implications for atherosclerotic plaque stability. *J Clin Invest.* 1996;98:2572–2579.
14. Fu X, Kassim SY, Parks WC, Heinecke JW. Hypochlorous acid oxygenates the cysteine switch domain of pro-matrilysin (MMP-7). A mechanism for matrix metalloproteinase activation and atherosclerotic plaque rupture by myeloperoxidase. *J Biol Chem.* 2001;276:41279–41287.
15. Yusuf S, Dagenais G, Pogue J, Bosch J, Sleight P. Vitamin E supplementation and cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *N Engl J Med.* 2000;342:154–160.
16. Stephens NG, Parsons A, Schofield PM, Kelly F, Cheeseman K, Mitchinson MJ. Randomised controlled trial of vitamin E in patients with coronary

- disease: Cambridge Heart Antioxidant Study (CHAOS) *Lancet*. 1996;347:781-786.
17. ACC/AHA Guidelines for the Management of Patients With ST-Elevation Myocardial Infarction- Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 1999 Guidelines for the Management of Patients With Acute Myocardial Infarction) *Circulation*, 2004;110:588-636.
  18. Braunwald E, Kloner RA. The stunned myocardium: prolonged, postischemic ventricular dysfunction. *Circulation*. 1982;66:1146-1149.
  19. Miller FC, Krucoff MW, Satler LF, Green CE, Fletcher RD, Del Negro AA, Pearle DL, Kent KM, Rackley CE. Ventricular arrhythmias during reperfusion. *Am Heart J*. 1986;112:928-932.
  20. Rezkalla SH, Kloner RA. No-reflow phenomenon. *Circulation*. 2002;105:656-662
  21. Verma S, Fedak PW, Weisel RD, Butany J, Rao V, Maitland A, Li RK, Dhillon B, Yau TM. Fundamentals of reperfusion injury for the clinical cardiologist. *Circulation*. 2002;105:2332-2336.
  22. Kloner RA. Does reperfusion injury exist in humans? *J Am Coll Cardiol*. 1993;21:537-545.
  23. Gross GJ, Kersten JR, Warltier DC. Mechanisms of postischemic contractile dysfunction. *Ann Thorac Surg*. 1999;68:1898- 1904.
  24. Bolli R, Jeroudi MO, Patel BS, et al. Direct evidence that oxygen-derived free radicals contribute to postischemic myocardial dysfunction in the intact dog. *Proc Natl Acad Sci U S A*. 1989;86:4695- 4699
  25. Jordan JE, Zhao ZQ, Vinten-Johansen J. The role of neutrophils in myocardial ischemia-reperfusion injury. *Cardiovasc Res*. 1999; 43:860-878.

26. Gross GJ, Farber NE, Hardman HF, Warltier DC. Beneficial actions of superoxide dismutase and catalase in stunned myocardium of dogs. *Am J Physiol.* 1986;250:H372-377.
27. Bolli R, Jeroudi MO, Patel BS, Aruoma OI, Halliwell B, Lai EK, McCay PB. Marked reduction of free radical generation and contractile dysfunction by antioxidant therapy begun at the time of reperfusion. Evidence that myocardial "stunning" is a manifestation of reperfusion injury. *Circ Res.* 1989;65:607-622.
28. Yamaguchi T, Shioji I, Sugimoto A, Komoda Y, Nakajima H. Chemical structure of a new family of bile pigments from human urine. *J Biochem.* 1994;116:298-303.
29. Izumi Y, Yamazaki M, Shimizu S, Shimizu K, Yamaguchi T, Nakajima H. Anti-bilirubin monoclonal antibody. II. Enzyme-linked immunosorbent assay for bilirubin fractions by combination of two monoclonal antibodies. *Biochim Biophys Acta.* 1988;967:261-266.
30. Shimomura H, Ogawa H, Takazoe K, Soejima H, Miyamoto S, Sakamoto T, Kawano H, Suefuji H, Nishikawa H, Arai H, Hokamaki J, Kajiwara I, Kugiyama K, Yoshimura M. Comparison of urinary biopyrrin levels in acute myocardial infarction (after reperfusion therapy) versus stable angina pectoris and their usefulness in predicting subsequent cardiac events. *Am J Cardiol.* 2002;90:108-111.
31. Pfeffer MA, Braunwald E. Ventricular remodeling after myocardial infarction: experimental observations and clinical implications. *Circulation.* 1990;81:1161- 1172.
32. Harada K, Sugaya T, Murakami K, Yazaki Y, Komuro I. Angiotensin II type 1A receptor knockout mice display less left ventricular remodeling and improved survival after myocardial infarction. *Circulation.* 1999;100: 2093- 2099.

33. Kinugawa S, Tsutsui H, Hayashidani S, et al. Treatment with dimethylthiourea prevents left ventricular remodeling and failure after experimental myocardial infarction in mice: role of oxidative stress. *Circ Res.* 2000;87:392- 398.
34. Siwik DA, Tzortzis JD, Pimental DR, et al. Inhibition of copper-zinc superoxide dismutase induces cell growth, hypertrophic phenotype, and apoptosis in neonatal rat cardiac myocytes in vitro. *Circ Res.* 1999;85: 147- 153.
35. Shiomi T, Tsutsui H, Matsusaka H, Murakami K, Hayashidani S, Ikeuchi M, Wen J, Kubota T, Utsumi H, Takeshita A. Overexpression of glutathione peroxidase prevents left ventricular remodeling and failure after myocardial infarction in mice. *Circulation.* 2004;109:544-9.
36. Nakamura K, Fushimi K, Kouchi H, Mihara K, Miyazaki M, Ohe T, Namba M. Inhibitory effects of antioxidants on neonatal rat cardiac myocyte hypertrophy induced by tumor necrosis factor-alpha and angiotensin II. *Circulation.* 1998;98:794-799.
37. Engberding N, Spiekermann S, Schaefer A, Heineke A, Wiencke A, Muller M, Fuchs M, Hilfiker-Kleiner D, Hornig B, Drexler H, Landmesser U. Allopurinol attenuates left ventricular remodeling and dysfunction after experimental myocardial infarction: a new action for an old drug? *Circulation.* 2004;110:2175-2179.
38. Ide T, Tsutsui H, Hayashidani S, Kang D, Suematsu N, Nakamura K, Utsumi H, Hamasaki N, Takeshita A. Mitochondrial DNA damage and dysfunction associated with oxidative stress in failing hearts after myocardial infarction. *Circ Res.* 2001;88:529-535.
39. Jun Hokamaki, Hiroaki Kawano, Michihiro Yoshimura, Hirofumi Soejima, Shinzo Miyamoto, Ichiro Kajiwara, Sunao Kojima, Tomohiro Sakamoto, Seigo Sugiyama, Nobutaka Hirai, Hideki Shimomura, Yasuhiro Nagayoshi, Kenichi

- Tsujita, Izuru Shioji, Shinya Sasaki, Hisao Ogawa. Urinary biopyrrins levels are elevated in relation to severity of heart failure. *J Am Coll Cardiol*. 2004;43:1880-1885.
40. Kasai H, Nishimura S. Hydroxylation of deoxyguanosine at the C-8 position by ascorbic acid and other reducing agents. *Nucleic Acids Res*. 1984;12:2137-2145.
  41. Shigenaga MK, Gimeno CJ, Ames BN. Urinary 8-hydroxy-2'-deoxyguanosine as a biological marker of in vivo oxidative DNA damage. *Proc Natl Acad Sci U S A*. 1989;86:9697-9701.
  42. Shibutani S, Takeshita M, Grollman AP. Insertion of specific bases during DNA synthesis past the oxidation-damaged base 8-oxodG. *Nature*. 1991;349(6308):431-434.
  43. Nakabeppu Y, Tsuchimoto D, Ichinoe A, Ohno M, Ide Y, Hirano S, Yoshimura D, Tominaga Y, Furuichi M, Sakumi K. Biological significance of the defense mechanisms against oxidative damage in nucleic acids caused by reactive oxygen species: from mitochondria to nuclei. *Ann N Y Acad Sci*. 2004;1011:101-111.
  44. Sakumi K, Tominaga Y, Furuichi M, Xu P, Tsuzuki T, Sekiguchi M, Nakabeppu Y. Ogg1 knockout-associated lung tumorigenesis and its suppression by Mth1 gene disruption. *Cancer Res*. 2003;63(5):902-905
  45. Wiseman H, Halliwell B. Damage to DNA by reactive oxygen and nitrogen species: role in inflammatory disease and progression to cancer. *Biochem J*. 1996;313:17-29.
  46. Tagesson C, Kallberg M, Klintenberg C, Starkhammar H. Determination of urinary 8-hydroxydeoxyguanosine by automated coupled-column high performance liquid chromatography: a powerful technique for assaying in vivo oxidative DNA damage in cancer patients. *Europe J Cancer*. 1995;31A:934-940.

47. Leinonen J, Lehtimäki T, Toyokuni S, Okada K, Tanaka T, Hiai H, Ochi H, Laippala P, Rantalaiho V, Wirta O, Pasternack A, Alho H. New biomarker evidence of oxidative DNA damage in patients with non-insulin-dependent diabetes mellitus. *FEBS Letters*. 1997;417:150–152.
48. Loft S, Vistisen K, Ewertz M, Tjønneland A, Overvad K, Poulsen HE. Oxidative DNA damage estimated by 8-hydroxydeoxyguanosine excretion in humans: influence of smoking, gender and body mass index. *Carcinogenesis*. 1992;13:2241–2247.
49. Martinet W, Knaapen MW, De Meyer GR, Herman AG, Kockx MM. Elevated levels of oxidative DNA damage and DNA repair enzymes in human atherosclerotic plaques. *Circulation*. 2002;106:927–932.
50. Nishikawa T, Sasahara T, Kiritoshi S, Sonoda K, Senokuchi T, Matsuo T, Kukidome D, Wake N, Matsumura T, Miyamura N, Sakakida M, Kishikawa H, Araki E. Evaluating of urinary 8-Hydroxydeoxyguanosine as a novel biomarker of macrovascular complications in Type 2 Diabetes. *Diabetes Care*. 2003;26:1507–1512.
51. The Thrombolysis in Myocardial Infarction (TIMI) trial. Phase I findings. TIMI Study Group. *N Engl J Med*. 1985;312:932–936.
52. McCord JM. Oxygen-derived free radicals in postischemic tissue injury. *New Engl J Med*. 1985;312:159–163.
53. Goldhaber JI, Weiss JN. Oxygen free radicals and cardiac reperfusion abnormalities. *Hypertension*. 1992;20:118–127.
54. Miyamoto S, Sakamoto T, Soejima H, Shimomura H, Kajiwara I, Kojima S, Hokamaki J, Sugiyama S, Yoshimura M, Ozaki Y, Nakamura H, Yodoi J, Ogawa H. Plasma thioredoxin levels and platelet aggregability in patients with acute myocardial infarction. *Am Heart J*. 2003;146:465–471.
55. Hokamaki J, Kawano H, Soejima H, Miyamoto S, Kajiwara I, Kojima S, Sakamoto T, Sugiyama S, Yoshimura M, Nakamura H, Yodoi J, Ogawa H. Plasma

- thioredoxin levels in patients with unstable angina. *Int J Cardiol.* 2005;99:225–231.
56. De Scheerder IK, van de Kraay AM, Lamers JM, Koster JF, de Jong JW, Serruys PW. Myocardial malondialdehyde and uric acid release after short-lasting coronary occlusions during coronary angioplasty: potential mechanisms for free radical generation. *Am J Cardiol.* 1991;68:392–395.
57. Ely D, Dunphy G, Dollwet H, Richter H, Sellke F, Azodi M. Maintenance of left ventricular function (90%) after twenty-four-hour heart preservation with deferoxamine. *Free Rad Biol Med.* 1992;12:479–485.
58. Young IS, Purvis JA, Lightbody JH, Adgey AA, Trimble ER. Lipid peroxidation and antioxidant status following thrombolytic therapy for acute myocardial infarction. *Eur Heart J.* 1993;14:1027–1033.
59. Cottin Y, Doise JM, Maupoil V, Tanniere-Zeller M, Dalloz F, Maynadie M, Walker MK, Louis P, Carli PM, Wolf JE, Rochette L. Plasma iron status and lipid peroxidation following thrombolytic therapy for acute myocardial infarction. *Fundam Clin Pharmacol.* 1998;12:236–241.
60. Miwa S, Toyokuni S, Nishina T, Nomoto T, Hiroyasu M, Nishimura K, Komeda M. Spatiotemporal alteration of 8-hydroxy-2'-deoxyguanosine levels in cardiomyocytes after myocardial infarction in rats. *Free Radical Research.* 2002;36:853–858.
61. Cordis GA, Maulik G, Bagchi D, Riedel W, Das DK. Detection of oxidative DNA damage to ischemic reperfused rat hearts by 8-hydroxydeoxyguanosine formation. *J Mol Cell Cardiol.* 1998;30:1939–1944.
62. Tsujita K, Shimomura H, Kawano H, Hokamaki J, Fukuda M, Yamashita T, Hida S, Nakamura Y, Nagayoshi Y, Sakamoto T, Yoshimura M, Arai H, Ogawa H. Effects of edaravone on reperfusion injury in patients with acute myocardial infarction. *Am J Cardiol.* 2004;94:481–484.

63. Watanabe T, Egawa M. Effects of an antistroke agent MCI-186 on cerebral arachidonate cascade. *J Pharmacol Exp Ther.* 1994;271:1624-1629.
64. Edaravone Acute Infarction Study Group. Effect of a novel free radical scavenger, edaravone (MCI-186), on acute brain infarction. Randomized, placebo-controlled, double-blind study at multicenters. *Cerebrovasc Dis.* 2003;15:222-229.
65. Li YY, Chen D, Watkins SC, Feldman AM. Mitochondrial abnormalities in tumor necrosis factor- $\alpha$ -induced heart failure are associated with impaired DNA repair activity. *Circulation.* 2001;104:2492-2497.
66. Karahalil B, Hogue BA, de Souza-Pinto NC, Bohr VA. Base excision repair capacity in mitochondria and nuclei: tissue-specific variations. *FASEB J.* 2002;16:1895-1902.
67. Tsutsui H, Ide T, Shiomi T, Kang D, Hayashidani S, Suematsu N, Wen J, Utsumi H, Hamasaki N, Takeshita A. 8-Oxo-dGTPase, which prevents oxidative stress-induced DNA damage, increases in the mitochondria from failing hearts *Circulation.* 2001;104:2883-2885.
68. Yasuhiro Nagayoshi, Jun Hokamaki, Hisao Ogawa, Kenichi Tsujita. Hideki Shimomura. Urinary 8-hydroxy-2'-deoxyguanosine to urinary creatinine ratio is correlated with severity of cardiac function in patients with old myocardial infarction. *Circulation.* 2004;110,17:III-704,